arXiv:1208.3419v1 [quant-ph] 16 Aug 2012

Equilibration of Measurement Statistics Under Complex Dynamics

Cozmin Ududec[1,2], Nathan Wiebe[2,3], and Joseph Emerson[2,4]
[1] *Department of Physics and Astronomy, University of Waterloo, Waterloo, ON, Canada*
[2] *Institute for Quantum Computing, 200 University Ave West, Waterloo, ON, Canada*
[3] *Department of Combinatics & Opt., University of Waterloo, Waterloo, ON, Canada and*
[4] *Department of Applied Math, University of Waterloo, Waterloo, ON, Canada*

There has been considerable recent progress in clarifying the conditions under which quantum systems equilibrate. We demonstrate that effective equilibration of an isolated system can occur under much weaker conditions than previously considered, by requiring only an assumption of sufficiently complex dynamics (which encodes a condition on the complexity of the system eigenvectors) and natural information-theoretic constraints resulting from the infeasibility of collecting an astronomically large amount of measurement data. In particular, we prove that, for generic Hamiltonian systems, the measurement statistics for maximally fine-grained measurements on pure states are effectively indistinguishable from the micro-canonical distribution after a finite equilibration time. We demonstrate that our assumptions correspond to physically realistic conditions by numerically confirming their validity for both generic many-body systems limited to two-body interactions and the quantum kicked-top in a globally chaotic regime.

Introduction. The mechanisms by which the characteristic features of quantum statistical mechanics emerge from an exact treatment of the quantum system are still not fully understood. In particular, it is not obvious how (or even whether) a pure state evolving according to some quantum dynamics approaches some form of equilibrium. Recently however, there has been a resurgence of interest in this problem [1–7], and progress in providing rigorous justification for the existence and time-scale of equilibration of small subsystems of a larger system [8–16].

Our work extends on this body of knowledge by showing that an assumption of complex dynamics is sufficient for an effective equilibration of measurement statistics for isolated quantum systems, without invoking mixed states, decoherence, subsystem traces or coarse-graining of observables. The form of equilibration we will study is an *information theoretic* one based on well-motivated operational constraints on the difficulty of predicting the detailed evolution of the system and the infeasibility of collecting an astronomically large amount of measurement data. We show that such equilibration is actually generic in the sense that it holds for almost all Hamiltonian systems, where "almost all" is defined with respect to the Gaussian Unitary Ensemble (GUE).

We further show that the strong form of randomness associated with GUE models is not necessary for microcanonical equilibration by identifying a weaker sufficient condition on the complexity of the system eigenvectors. We examine numerically whether this sufficient condition is met by – and effective information theoretic equilibration follows for – two important classes of physical systems: (i) generic many-body systems restricted to two-body interactions (2-local Hamiltonian systems), as well as (ii) single-body, chaotic dynamical systems in the macroscopic limit, in particular, the quantum kicked top [17] in a globally chaotic regime. This suggests that information-theoretic equilibration is not just mathematically generic, with respect to GUE, but may also be generic for more physically realistic models.

Our approach follows an established tradition of modelling complex physical systems with an appropriate ensemble of random matrices, a widely succesful approach that has been applied to many-body systems such as heavy nuclei and mesoscopic structures, as well as few-body but high-dimensional quantum systems associated with classically chaotic counterparts [17, 18]. This work develops and extends related ideas about the relevance of the randomness of the system's eigenvector statistics as a mechanism for microcanonical equilibration proposed earlier by Peres [19] and Srednicki [20, 21] and gives a theoretical explanation of the numerical equilibration results in Refs. [22, 23].

Setup. Consider a dynamically isolated quantum system described by a Hilbert space $\mathcal{H} = \mathbb{C}^D$, and which is in an initial state ρ_0. Further, let H be a fixed Hamiltonian describing the reversible evolution of the system, $\rho(t) = U(t)\rho_0 U(t)^\dagger$, where $U(t) = e^{-itH}$ (taking $\hbar = 1$). A quantum system is said to equilibrate if its state $\rho(t)$ evolves towards some particular mixed state σ_{eq} and remains close to it (i.e., remains indistinguishable from it in some practical sense [1, 9]) for almost all times. If there is such an equilibrium state, it must be the infinite time average state (sometimes called the generalized Gibbs state) $\sigma_\infty := \lim_{\tau\to\infty} \frac{1}{\tau}\int_0^\tau \rho(t)dt$. Note that, without further assumptions, σ_∞ may differ from the microcanonical state $\sigma_{mc} := \mathbb{1}/D$ or any thermal state [4]. The above form of equilibration has only been rigorously proven for the states of sufficiently small subsystems of the full system, and depends on the generic nature of random quantum states and their entanglement properties [5, 8–16]. Of course, finite dimensional closed quantum systems cannot equilibrate in this strong sense, as any finite dimensional unitary evolution is quasi-periodic.

In the following we instead consider a mathematically weaker but operationally well-motivated form of equilibration for the statistics of *global observables*. To this end, we consider any maximally fine-grained (non-degenerate) global observable A acting on $\mathcal{H}$, with spectral decomposition $A = \sum_{k=1}^{D} a_k \hat{P}_k$ and rank-one projectors $\hat{P}_k$. The primitive quantities of interest are the probabilities over outcome labels k,

$$\Pr\left(k|\rho(t)\right) = \mathrm{Tr}\left[\hat{P}_k U(t) \rho_0 U(t)^\dagger\right]. \tag{1}$$

Clearly, if we can show equilibration occurs even for such maximal observables then equilibration will occur also for more the kinds of coarse-grained observables, which provide even less information per observation.

Because we are interested in the dynamical properties of a probability distribution, we use an information theoretic criterion for equilibration based on the difficulty of distinguishing between the precise quantum distribution $\Pr\left(k|\rho(t)\right)$ and the micro-canonical distribution.

Definition 1. *A state $\rho(t)$ undergoing unitary dynamics $U(t)$ on $\mathcal{H} = \mathbb{C}^D$ equilibrates* information theoretically *with respect to an observable A, after a time t_{eq}, if for almost all times $t > t_{\mathrm{eq}}$ the outcome distribution $\Pr\left(k|\rho(t)\right)$ cannot be distinguished from the micro-canonical distribution $\Pr_{\mathrm{mc}}(k)$ with high probability with computational resources, including the number of samples taken from $\Pr\left(k|\rho(t)\right)$, that scale at most as $O(\mathrm{polylog}(D))$.*

This definition captures the idea that although the exact quantum distribution for the system may be *in principle* distinguishable from the micro-canonical distribution, the two are effectively indistinguishable if the resources needed to distinguish them exceed those practically available. We choose to delineate the practical from the impractical by requiring that the resources, and in particular the number of measurement outcomes, by disallowing resources that grow faster than poly–logarithmically with the Hilbert space dimension, because for many-body systems D scales exponentially with the number of subsystems. Of course, for a different physical scenario, a different cut-off may be appropriate. The important point is that, without the ability to efficiently predict $\Pr\left(k|\rho(t)\right)$, information theoretic equilibration occurs whenever the outcome variance,

$$\mathbb{V}_k\{\Pr\left(k\right)\} := D^{-1} \sum_k \left[\Pr\left(k\right) - D^{-1}\right]^2, \tag{2}$$

is sufficiently small. Note that, for simplicity of exposition, we are focussing on cases where the micro-canonical distribution is uniform $\Pr_{\mathrm{mc}}(k) := 1/D$.

Theorem 1. *Consider a system which generates random outcomes chosen from the set $\{1, \ldots, D\}$ according to an unknown distribution $\Pr(k)$ that is promised to have an outcome variance $\mathbb{V}_k\{\Pr(k)\}$ that scales as $O(D^{-2})$. Then at least $O(D^{1/4})$ samples from the distribution $\Pr\left(k\right)$ are needed in order to distinguish it from $\Pr_{mc}(k)$ with high probability.*

The proof is provided in the supplemental material. The key question is now: which classes of Hamiltonian systems satisfy the assumptions of Theorem 1 and hence exhibit information theoretic equilibration?

Generic Hamiltonians. In this section we show that information theoretic equilibration is mathematically generic in the sense that it holds for almost all Hamiltonian systems, where "almost all" is defined with respect to Gaussian Unitary Ensemble (GUE) [17, 18]. GUE is the unique unitarily invariant distribution over Hermitian matrices H that factorizes into a product of distributions each over an individual element of H. In particular, each element of H is an i.i.d. Gaussian random variable. More details can be found in Appendix B.

Our motivation for considering GUE is based on the insights from decades of experimental and numerical studies of complex many-body systems and few-body quantum chaos models in large Hilbert spaces [17, 18, 24]. The well-established *random matrix conjecture* states that certain experimentally relevant features of these complex systems, such as the spacings between adjacent eigenvalues, can be modeled statistically by the those of a random matrix drawn from an appropriate ensemble sharing the same symmetries.

In practice one often models the feature of interest with the ensemble mean, rather than a random sample from the ensemble. The feature of the random matrix ensemble that justifies this substitution is the phenomenon known as *concentration of measure*, which describes situations where the distribution of the property is tightly peaked about the ensemble mean. Concentration of measure plays a key role in our argument, and it is important for the reader to bear in mind that our averages over the ensemble are not an implicit form of decoherence or mixing, but an established technique for predicting the properties of indvidual systems. Indeed we now show that equilibration is generic by demonstrating that the concentration of measure phenomena holds even for the outcome statistics of maximal observables.

Theorem 2. *Take a non-degenerate observable A acting on $\mathcal{H} = \mathbb{C}^D$, an initial pure state $\rho_0 = |x\rangle\langle x|$ which is an eigenstate of A, and an unknown unitary $U(t) = e^{-itH}$ where H is drawn uniformly at random from GUE. Then there exists a $t_{\mathrm{eq}}(D) = O(D^{-1/6})$ s. t. for all $t > t_{\mathrm{eq}}(D) = O(D^{-1/6})$ the mean, variance, and fourth moment of the measurement outcome probabilities $\Pr\left(k|x,t\right)$ over GUE are given by:*

$$\mathbb{E}_{\mathrm{GUE}}\{\Pr\left(k|x,t\right)\} = \frac{D + \delta_{xk} O(D)}{D^2 - 1} + O(D^{-2}), \tag{3}$$

$$\mathbb{V}_{\mathrm{GUE}}\{\Pr\left(k|x,t\right)\} = O(D^{-2}), \tag{4}$$

$$\mathbb{E}_{\mathrm{GUE}}\left\{\Pr\left(k|x,t\right)^4\right\} = O(D^{-4}). \tag{5}$$

For a proof see the supplemental material. This result implies that $\Pr\left(k|x,t\right)$ is concentrated around $1/D$ for almost every Hamiltonian chosen from the GUE and $t > t_{\mathrm{eq}}(D)$. In particular, by Chebyshev's inequality, in the limit of large D, with high probability over H drawn from GUE, the corresponding probabilities $\Pr\left(k|x,t\right)$ are at most a constant multiple of $1/D$:

$$\Pr_{H\sim\mu_D}\left(\left|\Pr(k|x,t)-\mathbb{E}_{\mathrm{GUE}}\{\Pr(k|x,t)\}\right|\geq\epsilon\right)\leq\frac{O(D^{-2})}{\epsilon^2}, \tag{6}$$

where μ_D is the appropriate measure for the GUE. In other words, the probabilities $\Pr(k|x,t)$ for an individual system are very close to the ensemble average.

These results imply a concentration of the outcome variance. Expanding the outcome and GUE variances and using Theorem 2, we find that, for $t > t_{\mathrm{eq}}(D)$,

$$\begin{aligned}&\mathbb{V}_{\mathrm{GUE}}\{\mathbb{V}_k\{\Pr(k|x,t)\}\} \qquad (7)\\ =\ &D^{-2}\sum_{j,k}\mathbb{E}_{\mathrm{GUE}}\left\{\Pr(k|x,t)^2\Pr(j|x,t)^2\right\}+O(D^{-4}).\end{aligned}$$

Then using the Cauchy-Schwartz inequality for expectations and Theorem 2 again, we have:

$$\sum_{j,k}\mathbb{E}_{\mathrm{GUE}}\left\{\Pr(k|x,t)^2\Pr(j|x,t)^2\right\}\leq O(D^{-2}). \tag{8}$$

This proves that $\mathbb{V}_{\mathrm{GUE}}\{\mathbb{V}_k\{\Pr(k|x,t)\}\} = O(D^{-4})$. Chebyshev's inequality then implies (in the same fashion as (6)) that $\mathbb{V}_k\{\Pr(k|x,t)\}\in O(D^{-2})$ with high probability over GUE.

Theorem 1 therefore implies that $\Pr(k|x,t)$ cannot be efficiently distinguished from $\Pr_{\mathrm{mc}}(k)$. Hence, in high–dimensional Hilbert spaces, the vast majority of *individual* Hamiltonian systems (which are at all times in a pure state), equilibrate information-theoretically. These results imply, rather remarkably, that the dynamical pure-state quantum fluctuations for $t > t_{eq}$ become vanishingly small (in an experimentally relevant sense) in the limit of large D.

Sufficient conditions for Theorems 1 and 2. It is interesting to note that the very strong assumptions of randomness implicit in GUE (specifically, the Haar randomness of the energy eigenstates) are not necessary for Theorems 1 and 2 to hold, and identifying weaker sufficient conditions is desirable.

As shown in the supplemental material, the object that underlies the distribution of $\Pr(k|x,t)$ is the unitary, C, which transforms the eigenbasis of H to that of the observable A. Working in the eigenbasis of A, we can write $U(t) = CF(t)C^\dagger$, where $F(t) = \mathrm{diag}[e^{-itE_a}]$, and $\{E_a\}_{a=1}^D$ are the energy eigenvalues of H. Given an initial pure state $|x\rangle\langle x|$, the measurement outcome probabilities can then be written as

$$\Pr(k|x,t)=\mathrm{Tr}\left[|k\rangle\langle k|CF(t)C^\dagger|x\rangle\langle x|CF(t)^\dagger C^\dagger\right]. \tag{9}$$

Information theoretic equilibration requires that $\mathbb{V}_k\{\Pr(k|x,t)\}\in O(D^{-2})$, which in turn requires that we know certain properties of $\Pr(k|x,C,F(t))^2$. It is not difficult to see that $\Pr(k|x,t)$ and $\Pr(k|x,t)^2$ can be concisely represented by

$$\Pr(k|x,t)=\langle L_2|C^{\otimes 2}\otimes\bar{C}^{\otimes 2}|R_2(t)\rangle, \tag{10}$$

$$\Pr(k|x,t)^2=\langle L_4|C^{\otimes 4}\otimes\bar{C}^{\otimes 4}|R_4(t)\rangle, \tag{11}$$



FIG. 1: The average over x of the RLH ensemble average of the probabilities $P(x\neq 0|0,t)$ of the evolved state plotted as a function of time for 5–, . . . , 10–qubit systems. This average equilibrates to $1/(D+1)$ within an equilibration time that scales approximately as $O(1/\sqrt{\log(D)})$

.

where $\langle L_2| = \langle k,x,x,k|$, $\langle L_4| = \langle k,x,k,x,x,k,x,k|$, and

$$|R_2(t)\rangle=\sum_{b,b'}e^{it(E_b-E_{b'})}|b,b',b,b'\rangle, \tag{12}$$

$$|R_4(t)\rangle=\sum_{b,b',d,d'}e^{it(E_b-E_{b'}+E_d-E_{d'})}|b,b',d,d',b,b',d,d'\rangle. \tag{13}$$

We refer to a term of the form $\langle L_2|C^{\otimes 2}\otimes\bar{C}^{\otimes 2}|b,b',b,b'\rangle$ in the sum in (10) as a $(2,2)$–term because there are two basis change matrices acting on each factor space. The analogous terms in (11) will be called $(4,4)$–terms.

We see from the above that the strong randomness of GUE eigenvectors is not necessary, and that scalings given in eqns. (3),(4) and (5) are satisfied provided that the matrix elements of C, namely the Hamiltonian eigenvector components, satisfy a unitary t-design condition [25]. This insight has been identified recently in the context of subsystem equilibration in Refs. [10, 11, 16], where one requires eigenvector component reflecting GUE-randomness only for polynomials of degree at most $t = 4$. For microcanonical equilibration our proof technique shows that eigenvector components reflecting GUE-randomness for polynomials of degree at most $t = 8$ is sufficient to imply that information theoretic equilibration is typical using our Theorems 1 and 2. We now provide numerical evidence that such scalings are realized for typical systems drawn from a more physically realistic ensemble consisting of 2–local Hamiltonians.

Equilibration of Local Hamiltonians. The class of k–local Hamiltonians consists of those Hamiltonians composed of at most k–body interactions (for fixed k) between d–level quantum systems. These Hamiltonians are ubiquitous in condensed matter theory and quantum computing [26]. We anticipate that such Hamiltonians have sufficient complexity to induce equilibration because the



FIG. 2: Numerically computed expectation values over k and b of the RLH ensemble average and variances of the dominant $(4,4)$ term, $\langle x,k,k,x|C^{\otimes 2}\otimes\bar{C}^{\otimes 2}|b,b,b,b\rangle$, plotted as a function of $D=2^n$, and compared to the analytically computed scalings for GUE. The variance of the $(4,4)$ terms scale as $3790/D^8$ for $D>128$.



FIG. 3: The measurement outcome variance $\mathbb{V}_k\{\Pr(k|x)\}$ plotted as a function of system size, both before (2 kicks) and after (10 kicks) the equilibration time. For 10 kicks, we see excellent agreement with the GUE prediction of $O(D^{-2})$, confirming information theoretic equilibration of a one-body system in the macroscopic limit in a regime of global chaos ($\boldsymbol{\alpha}=[1.1,1.0,1.0]$ and $\boldsymbol{\tau}=\mathrm{diag}(10,0,1)$).

interactions they are composed of generate a universal gate set for quantum computing. Although difficult to prove analytically, we provide numerical evidence that almost all randomly chosen 2–local Hamiltonians acting on qubits equilibrate information theoretically with respect to the observable $A=\sigma_z^1\otimes\cdots\otimes\sigma_z^n$ (corresponding to a non-degenerate projective measurement in the computational basis). This describes a wide-class of global observables because the ensemble of random 2–local Hamiltonians is invariant with respect to independent rotations of the individual qubits.

We construct the ensemble of random 2–local Hamiltonians on n systems (denoted RLH) as follows:

$$H=\sum_{i=1}^{n}\sum_{p}a_i\sigma_p^{(i)}+\sum_{i\neq j}\sum_{p}\sum_{p'}b_{i,j}\sigma_p^{(i)}\sigma_{p'}^{(j)}, \tag{14}$$

where $p,p'\in\{X,Y,Z\}$, and each a_i and $b_{i,j}$ is a Gaussian random variable with mean 0 and variance 1.

Fig. 1 shows that systems evolving under RLH Hamiltonians approach equilibrium as D increases. The equilibration time can be difficult to define, but for concreteness we estimate it using the location of the inflection points of the curves in Fig. 1. These times scale as $O(1/\sqrt{\log(D)})$, similar to what would be expected from analysis of the Ehrenfest breaktime for quantum chaotic systems [17, 22, 23]. Fig. 2 further shows that the RLH average of the $(4,4)$–terms agrees with the GUE predictions of $O(D^{-4})$ scaling. We also find that the variance of the $(4,4)$ terms is $O(D^{-8})$ suggesting that concentration of measure holds for these. This implies that similarly to the GUE case, $\mathbb{V}_k\{\Pr(k|x,t)\}\in O(D^{-2})$ for an individual system, and hence the conditions of Theorem 1 are met for almost all H taken from the RLH provided that $t>t_{\mathrm{eq}}(D)=O(1/\sqrt{\log(D)})$. This is strong evidence that generic 2–local Hamiltonians exhibit information theoretic equilibration.

Equilibration for Individual Systems. Information theoretic equilibration of pure states can occur also for an individual system consisting of a one-body dynamical model associated with global classical chaos. A variant of the quantum kicked top, exhibits this property. It is described by the Floquet map (with $\hbar=1$) [17]:

$$U_F=e^{-i\boldsymbol{J}\boldsymbol{\tau}\boldsymbol{J}^T}e^{-i\mathbf{J}\cdot\alpha}, \tag{15}$$

where $\mathbf{J}=[J_x,J_y,J_z]$ is a vector of angular momentum operators, τ is the moment of inertia tensor, which is diagonal with entries $[\tau_{xx},\tau_{yy},\tau_{zz}]$ and $\alpha=[\alpha_{\mathbf{x}},\alpha_{\mathbf{y}},\alpha_{\mathbf{z}}]$ is a vector of kick–strengths. The dimension of the system is $D=2j+1$ where j is the total angular momentum quantum number. We see in Fig. 3 that for a set of chaotic parameters for λ and τ, roughly 10 kicks (applications of U_F) are needed in order for $\mathbb{V}_k\{\Pr(k|x,t)\}=O(D^{-2})$, and hence for Theorem 1 to apply.

Discussion. We have shown analytically that information theoretic equilibration holds generically for the probability distributions of maximal measurements of closed Hamiltonian systems that satisfy a condition of sufficient complexity. Our condition of sufficient complexity is a natural extension of the random matrix conjecture [27] to low-degree polynomials of the eigenvector components of complex quantum systems [28, 29]. We have also demonstrated numerically that this condition is satisfied by a class of random 2–local (many-body) Hamiltonians, as well as an individual single-body dynamical system in a regime of global classical chaos. Remarkably, information theoretic equilibration is observed to hold for pure quantum states for each of these systems, without requiring any form of decoherence or restricting to coarse-grained measurements.

In the context of random matrix models we showed that this weak form of equilibration holds for almost all Hamiltonians for any fixed observable. In the context of a fixed Hamiltonian model, one can ask: for which observables is this form of equilibration expected to hold? The answer is already clear: almost all of them, because the relevant feature is the *relative randomness* of the basis transformation from the energy eigenbasis to the observable's eigenbasis, or equivalently, the matrix of energy eigenvector components in the basis of the observable.

Our key insight is that although dynamical pure states exhibit coherent fluctuations away from true micro-canonical equilibrium, these fluctuations remain small and appear pseudo-random, and hence their detection requires extraordinary experimental resources, such as collecting $O(D^{1/4})$ measurement outcomes from repetitions of the exact experimental conditions, or pre–computation of the location of the dynamical state in a D-dimensional Hilbert space, or performing joint (entangling) measurements on identical copies of the system. In the absence of such resources, by Theorem 1 we see that after some finite time (which depends on the details of the energy spectrum) the probability distributions for dynamical pure states cannot be distinguished from the micro–canonical distribution.

Acknowledgements. We thank Fernando Brandao, Carl Caves, Patrick Hayden, Daniel Gottesman, and Victor Veitch for insightful comments. J. Emerson and C. Ududec acknowledge funding from CIFAR and NSERC, and thank the Perimeter Institute for Theoretical Physics where this work was brought to completion. Nathan Wiebe acknowledges funding from the US ARO/DTO and CIFAR.

[1] S. Popescu, A. J. Short, and A. Winter, Nature Physics **2**, 754 (2006), arXiv:0511225.

[2] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì (2010), arXiv:1003.2129.

[3] P. Reimann, New Journal of Physics **12**, 055027 (2010).

[4] M. Rigol, V. Dunjko, and M. Olshanii, Nature **452**, 854 (2008).

[5] A. J. Short and F. T. C., New Journal of Physics **14**, 013063 (2012), arXiv:1110.5759.

[6] C. L. Venuti and P. Zanardi, Phys. Rev. A **81**, 022113 (2010).

[7] M. Cramer and J. Eisert, New Journal of Physics **12**, 055020 (2010), arXiv:0911.2475.

[8] N. Linden, S. Popescu, A. J. Short, and A. Winter, Phys. Rev. E **79**, 061103 (2009).

[9] A. J. Short, New Journal of Physics **13**, 053009 (2011).

[10] F. Brandao, P. Cwiklinski, M. Horodecki, P. Horodecki, J. Korbicz, and M. Mozrzymas (2011), arXiv:1108.2985.

[11] L. Masanes, A. J. Roncaglia, and A. Acin (2011), arXiv:1108.0374.

[12] S. Lloyd, Ph.D. thesis, Rockefeller University, New York (1988).

[13] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì, Phys. Rev. Lett. **96**, 050403 (2006).

[14] J. Gemmer, M. Michel, and G. Mahler, *Quantum Thermodynamics: Emergence of Thermodynamic Behavior within Composite Quantum Systems.* (Springer: Berlin, 2004).

[15] A. Riera, C. Gogolin, and J. Eisert (2011).

[16] M. Znidaric, C. Pineda, and I. García-Mata, Phys. Rev. Lett. **107**, 080404 (2011), arXiv:1104.5263.

[17] F. Haake, *Quantum Signatures of Chaos* 2^{nd} *Edition* (Springer-Verlag New York, 2004).

[18] M. L. Mehta, *Random Matrices; 3rd ed.* (Elsevier, San Diego, CA, 2004).

[19] A. Peres, Phys. Rev. A **30**, 504 (1984).

[20] M. Srednicki, Phys. Rev. E **50**, 888 (1994).

[21] M. Srednicki (1994), arXiv:9410046.

[22] J. Emerson and L. Ballentine, Phys. Rev. A **63**, 052103 (2001).

[23] J. Emerson and L. E. Ballentine, Phys. Rev. E **64**, 026217 (2001).

[24] T. Guhr, A. Müller-Groeling, and H. A. Weidenmüller, Physics Reports **299**, 189 (1998).

[25] C. Dankert, R. Cleve, J. Emerson, and E. Livine, Phys. Rev. A **80**, 012304 (2009), arXiv:0606161.

[26] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge U.K., 2000).

[27] O. Bohigas, M. J. Giannoni, and C. Schmit, Phys. Rev. Lett. **52**, 1 (1984).

[28] M. Kus and K. Zyzckowski, Phys. Rev. A **44**, 956 (1991).

[29] M. J. Kus, M. and F. Haake, J. Phys. A: Math. Gen. **21**, L1073 (1988).

[30] R. Balian, Il Nuovo Cimento B (1965-1970) **57**, 183 (1968).

Appendix A: Proof of Theorem 1

Proof of Theorem 1. First, notice that the lack of knowledge of the form of the distribution $\Pr(k)$ implies that the labels of the particular outputs k are of no significance. This means that we can only use the frequency of coincidences in the output in order to distinguish $\Pr(k)$ from the uniform distribution.

It is straightforward to see that $O(\sqrt{D})$ observations are needed to be able to obtain a pair of identical measurement outcomes drawn from the uniform distribution. We compare this to a distribution with $\mathbb{V}_k\{\Pr(k)\} = O(D^{-2})$ that has the highest probability of coincidence for some outputs. We achieve this by imposing a restriction on $\Pr(k)$ that is weaker than $\mathbb{V}_k\{\Pr(k)\} = O(D^{-2})$ but is also implied by the variance constraint. Using the the definition of variance, $\mathbb{V}_k\{\Pr(k)\} = D^{-1}\sum_k(\Pr(k) - D^{-1})^2$, and expanding the sum and using $\sum_k \Pr(k) = 1$, we find

$$\mathbb{V}_k\{\Pr(k)\} = O(D^{-2}) \ \Rightarrow \Pr(k) \le O(D^{-1/2}),\ \forall\ k. \tag{A1}$$

We will now drop the variance restriction and find a distribution $\mathrm{Q}(k)$ that only satisfies $\mathrm{Q}(k) \le O(D^{-1/2})$ and also maximizes the probability of coincidence. This distribution is, for some constant a, given by

$$\mathrm{Q}(k) = \begin{cases} \sqrt{\frac{a}{D}} & : \ k \le \sqrt{\frac{D}{a}} \\ 0 & : \ \text{else} \end{cases}. \tag{A2}$$

Given the distribution Q, the probability that there are no coincidental outcomes after N samples is given by:

$$\frac{\left(\sqrt{\frac{D}{a}}\right)!}{\left(\sqrt{\frac{D}{a}}-N+1\right)!\left(\sqrt{\frac{D}{a}}\right)^{N}} = 1-\frac{N(N-1)}{2\sqrt{\frac{D}{a}}}+O(N/D). \tag{A3}$$

Therefore at least $N = O(D^{1/4})$ samples are needed to conclude with high probability that $Q(k)$ is not uniform if $\mathrm{Q}(k) \leq O(D^{-1/2})$. This implies that a distribution $\Pr(k)$ with $\mathbb{V}_k\{\Pr(k)\} = O(D^{-2})$ cannot be distinguished from the uniform distribution with high probability unless $N \geq O(D^{1/4})$, as claimed. $\square$

Appendix B: Proof of Theorem 2

We prove Theorem 2 in two steps:

1. We derive expressions for the Haar expectations of $\Pr(k|x, C, F(t))$ (see (B2) below) and the second and fourth moments. See Lemmas 1, 2, and 3.
2. We find the expectations of these expressions over the GUE eigenvalue distribution, and show that there exists a time $t_{\mathrm{eq}}(D) = O(D^{-1/6})$ such that for all $t > t_{\mathrm{eq}}(D)$, eqns. (3), (4), and (5) hold. See Section B 2 for these expectations.

Since the calculations in this subsection require in depth knowledge of the properties of the Gaussian Unitary Ensemble (GUE), we will begin by giving a brief review of its properties [17, 18, 30]. A Hermitian matrix H generated according to the GUE has diagonal elements h_{aa} that are real valued random variables each with distribution $\mathcal{N}(0, \sigma^2)$, and off-diagonal elements h_{ab} with real and imaginary parts that are random variables each with distribution $\mathcal{N}(0, \frac{1}{2}\sigma^2)$. The variance σ^2 is a free parameter, which is closely related to the expected maximum energy eigenvalue as well as the ensemble average energy level spacing.

There are several important drawbacks to using GUE which deserve mention before moving on. First, the average level density, which takes the form of a semi-circle, is not a good model for physically relevant Hamiltonian systems, even chaotic ones. In particular, for most natural systems, such as those with only two particle interactions, the norm of the Hamiltonian scales polynomially with the number of particles [8]. However, the expected norm of a GUE Hamiltonian scales polynomially in the Hilbert space dimension D [17]. As we will see below, this has a large impact on what might be called the equilibration time for these dynamical systems, which should therefore be taken with a grain of salt. Because of this, and in order to simplify calculations, we will follow the standard practice [11, 18] of taking the variance $\sigma^2 = \frac{1}{2}$.

The joint distribution over all elements of H factorizes into a product of a distribution over eigenvectors, and a distribution over the energy eigenvalues of H (see [18] Theorem 3.3.1, or [17] Chapter 4). Further, the joint probability distribution over eigenvectors of H is the same as that over the change of basis matrix C, namely the Haar measure on the unitary group $\mathbb{U}(D)$ [17, 18]. Letting C be the unitary which takes the eigenbasis of H to that of A, and working in the eigenbasis of A, we can write

$$U(t) = CF(t)C^{\dagger}, \tag{B1}$$

where $F(t) := \mathrm{diag}[e^{-itE_a}]$, and $\{E_a\}_{a=1}^{D}$ are the energy eigenvalues of H. We can therefore take separate expectations over eigenvectors and eigenvalues, namely, over the matrices C and F. $\Pr(k|x,t)$ can be expressed as

$$\Pr(k|x, C, F(t)) = \mathrm{Tr}\left[|k\rangle\langle k|CF(t)C^{\dagger}|x\rangle\langle x|CF(t)^{\dagger}C^{\dagger}\right]. \tag{B2}$$

In the following we will write the expectation over the Haar measure on the unitary group $\mathbb{U}(D)$ of change of basis matrices C, as $\mathbb{E}_C\{\cdot\}$, and $\mathbb{E}_{\mathrm{spec}}\{\cdot\}$ for the expectation over the GUE eigenvalue distribution.

1. Expectations over eigenvectors

Lemma 1. *Take a non-degenerate observable A acting on $\mathcal{H} = \mathbb{C}^D$, an initial pure state $\rho_0 = |x\rangle\langle x|$ which is an eigenstate of A, and a unitary $U(t) = e^{-itH}$ where H is drawn uniformly at random from GUE. Then the variance of the measurement outcome probabilities over the Haar measure on the unitary group $\mathbb{U}(D)$ of change of basis matrices C is given by:*

$$\begin{aligned}
&\mathbb{V}_C\{\Pr(k|x, C, F(t))\} \\
&\quad= \frac{1}{D^4}\Big\{D^2 - 2D + 4 + (7-2D)\delta_{xk} \\
&\quad+ |\mu(t)|^2(2D\delta_{xk} - 10\delta_{xk} - 2) \\
&\quad+ \delta_{xk}|\mu(2t)|^2 + 2\delta_{xk}Re[\mu(t)^2\mu(-2t)]\Big\} + O(D^{-5}),
\end{aligned} \tag{B3}$$

where we have defined $\mu(t) = \mathrm{Tr}[U(t)] = \mathrm{Tr}[F(t)]$ (this is often called the spectral form factor [17, 18]).

Proof. First, note that the squares of the outcome probabilities $\Pr(k|x, C, F(t))$ can be written in the form:

$$\begin{aligned}
&\mathrm{Tr}\left[|k\rangle\langle k|CF(t)C^{\dagger}|x\rangle\langle x|CF(t)^{\dagger}C^{\dagger}\right]^2 \\
&\quad= \sum_{s,s'}\langle s|k\rangle\langle k|CF(t)C^{\dagger}|x\rangle\langle x|CF(t)^{\dagger}C^{\dagger}|s\rangle \\
&\quad\times\ \langle s'|k\rangle\langle k|CF(t)C^{\dagger}|x\rangle\langle x|CF(t)^{\dagger}C^{\dagger}|s'\rangle \\
&\quad= \langle L_4|C^{\otimes 4}\otimes\bar{C}^{\otimes 4}|R_4(t)\rangle,
\end{aligned} \tag{B4}$$

where $\bar{C}$ is the complex conjugate of C and

$$\begin{aligned}
\langle L_4| &= \langle k, x, k, x, x, k, x, k|, \\
|R_4(t)\rangle &= \sum_{b,b',d,d'} e^{it(E_b - E_{b'} + E_d - E_{d'})} \\
&\quad\times|b, b', d, d', b, b', d, d'\rangle,
\end{aligned} \tag{B5}$$

The expectation of the expression $C^{\otimes 4} \otimes \bar{C}^{\otimes 4}$ over Haar measure can be written as the projector onto the subspace spanned by the vectors

$$|\Phi_\pi\rangle = (V_\pi \otimes \mathbb{1})\,|\phi\rangle_{1,5}|\phi\rangle_{2,6}|\phi\rangle_{3,7}|\phi\rangle_{4,8}, \tag{B6}$$

where $|\phi\rangle_{ij} = \sum_{a=1}^{D} |a\rangle_i|a\rangle_j$, and the index π runs over the 4! permutations of the elements $\{1,2,3,4\}$, and V_π is the unitary permutting the first four factor spaces according to π. It was shown in [10] that this projector is given by

$$\mathbb{E}_C\{C^{\otimes 4} \otimes \bar{C}^{\otimes 4}\} = \sum_{\pi,\sigma} (M^{-1})_{\pi,\sigma} |\Phi_\pi\rangle\langle\Phi_\sigma|, \tag{B7}$$

where the matrix M has components $M_{\pi,\sigma} = \langle V_\pi|V_\sigma\rangle = \mathrm{Tr}[V_{\pi^{-1}}V_\sigma] = d^{l(\pi^{-1}\sigma)}$, and $l(\sigma)$ is number of cycles in the cycle decomposition of the permutation $\pi^{-1}\sigma$. We then have

$$\begin{aligned}&\mathbb{E}_C\Big\{\Pr(k|x,C,F(t))^2\Big\}\\ &\qquad = \sum_{\pi,\sigma} \langle L_4|\Phi_\pi\rangle (M^{-1})_{\pi,\sigma} \langle\Phi_\sigma|R_4(t)\rangle,\end{aligned} \tag{B8}$$

where the inner products $\langle\Phi_\sigma|R_4(t)\rangle$ are given by:

$$\begin{aligned}
\langle\Phi_{(1,2,3,4)}|R_4(t)\rangle &= |\mu(t)|^4,\\
\langle\Phi_{(1,2,4,3)}|R_4(t)\rangle = \langle\Phi_{(1,3,2,4)}|R_4(t)\rangle &= d|\mu(t)|^2,\\
\langle\Phi_{(2,1,3,4)}|R_4(t)\rangle = \langle\Phi_{(4,2,3,1)}|R_4(t)\rangle &= d|\mu(t)|^2,\\
\langle\Phi_{(1,3,4,2)}|R_4(t)\rangle = \langle\Phi_{(1,4,2,3)}|R_4(t)\rangle &= |\mu(t)|^2,\\
\langle\Phi_{(2,3,1,4)}|R_4(t)\rangle = \langle\Phi_{(2,4,3,1)}|R_4(t)\rangle &= |\mu(t)|^2,\\
\langle\Phi_{(3,1,2,4)}|R_4(t)\rangle = \langle\Phi_{(3,2,4,1)}|R_4(t)\rangle &= |\mu(t)|^2,\\
\langle\Phi_{(4,1,3,2)}|R_4(t)\rangle = \langle\Phi_{(4,2,1,3)}|R_4(t)\rangle &= |\mu(t)|^2,\\
\langle\Phi_{(3,4,1,2)}|R_4(t)\rangle &= |\mu(2t)|^2,\\
\langle\Phi_{(1,4,3,2)}|R_4(t)\rangle &= \mu(t)^2\bar{\mu}(2t),\\
\langle\Phi_{(3,2,1,4)}|R_4(t)\rangle &= \mu(2t)\bar{\mu}(t)^2,\\
\langle\Phi_{(2,1,4,3)}|R_4(t)\rangle = \langle\Phi_{(4,3,2,1)}|R_4(t)\rangle &= d^2,\\
\langle\Phi_{(2,3,4,1)}|R_4(t)\rangle = \langle\Phi_{(3,4,2,1)}|R_4(t)\rangle &= d,\\
\langle\Phi_{(2,4,1,3)}|R_4(t)\rangle = \langle\Phi_{(3,1,4,2)}|R_4(t)\rangle &= d,\\
\langle\Phi_{(4,1,2,3)}|R_4(t)\rangle = \langle\Phi_{(4,3,1,2)}|R_4(t)\rangle &= d,
\end{aligned}$$

Further, $\langle L_4|\Phi_\pi\rangle = 1$ for $\pi = (4,3,2,1)$, $(4,1,2,3)$, $(2,3,4,1)$, $(2,1,4,3)$, and $\langle L_4|\Phi_\pi\rangle = \delta_{ik}$ for all other π.

Taking the sum of the above terms with M^{-1} in eqn. (B8), we find:

$$\begin{aligned}&\mathbb{E}_C\Big\{\Pr(k|x,C,F(t))^2\Big\}\\
&= \frac{1}{\alpha}\Big\{|\mu(t)|^4\left((D^2-D-2)\delta_{xk}+2\right)\\
&+ |\mu(t)|^2\left(-4D^2-12D-8+(4D^3+8D^2+4D+8)\delta_{xk}\right)\\
&+ (D^2-D-2)\delta_{xk}\left(|\mu(2t)|^2+\bar{\mu}(2t)\mu(t)^2+\mu(2t)\bar{\mu}(t)^2\right)\\
&+ 2D^4+8D^3+6D^2-(4D^3+12D^2)\delta_{xk}\\
&+ 2|\mu(2t)|^2+2\bar{\mu}(2t)\mu(t)^2+2\mu(2t)\bar{\mu}(t)^2\Big\},\end{aligned} \tag{B9}$$

where $\alpha = D^2(D-1)(D+1)(D+2)(D+3)$. Using $\mathbb{E}_C\{\Pr(k|x,C,F(t))\}$ from Lemma 2, we then have

$$\begin{aligned}&\mathbb{V}_C\{\Pr(k|x,C,F(t))\}\\
&= \frac{1}{D^4}\Big\{D^2-2D+4+(7-2D)\delta_{xk}\\
&+ |\mu(t)|^2(2D\delta_{xk}-10\delta_{xk}-2)\\
&+ \delta_{xk}|\mu(2t)|^2+2\delta_{xk}Re[\mu(t)^2\mu(-2t)]\Big\}+O(D^{-5}).\end{aligned} \tag{B10}$$

□

Lemma 2. *Take a non-degenerate observable A acting on $\mathcal{H} = \mathbb{C}^D$, an initial pure state $\rho_0 = |x\rangle\langle x|$ which is an eigenstate of A, and a unitary $U(t) = e^{-itH}$ where H is drawn uniformly at random from GUE. Then the expectation of the measurement outcome probabilities for C distributed according to the Haar measure on the unitary group $\mathbb{U}(D)$ is given by:*

$$\mathbb{E}_C\{\Pr(k|x,C,F(t))\} = \frac{D-\delta_{xk}+|\mu(t)|^2\left(\delta_{xk}-\frac{1}{D}\right)}{D^2-1}. \tag{B11}$$

Proof. This expectation can be calculated in a similar but simpler fashion as the variance in the previous lemma, so we leave the proof as an exercise. □

Lemma 3. *Take a non-degenerate observable A acting on $\mathcal{H} = \mathbb{C}^D$, an initial pure state $\rho_0 = |x\rangle\langle x|$ which is an eigenstate of A, and a unitary $U(t) = e^{-itH}$ where H is drawn uniformly at random from the GUE. Then the fourth moment of the measurement outcome probabilities over the Haar measure on the unitary group $\mathbb{U}(D)$ of change of basis matrices C is given by:*

$$\begin{aligned}&\mathbb{E}_C\Big\{\Pr(k|x,C,F(t))^4\Big\}\\
&\quad \le \frac{1}{D^8}\sum_{\pi}\langle\Phi_\pi|R_8(t)\rangle + \sum_{\pi,\sigma} B_{\pi,\sigma}\langle\Phi_\sigma|R_8(t)\rangle,\end{aligned} \tag{B12}$$

where B is a matrix with components $\le O(D^{-9})$.

Proof. The fourth power of the outcome probabilities can be written in the form:

$$\begin{aligned}&\mathrm{Tr}\left[|k\rangle\langle k|CF(t)C^\dagger|x\rangle\langle x|CF(t)^\dagger C^\dagger\right]^4\\
&\qquad = \langle L_8|C^{\otimes 8}\otimes\bar{C}^{\otimes 8}|R_8(t)\rangle,\end{aligned} \tag{B13}$$

where $\langle L_8|$ and $|R_8(t)\rangle$ are defined analogously to (B5), but with twice the number of tensor factors. Further, the average of the expression $C^{\otimes 8} \otimes \bar{C}^{\otimes 8}$ over Haar measure can be written as the projector onto the subspace spanned by the vectors

$$|\Phi_\pi\rangle = (V_\pi \otimes \mathbb{1})\,|\phi\rangle_{1,9}|\phi\rangle_{2,10}\ldots|\phi\rangle_{8,16}, \tag{B14}$$

where the index π runs over the 8! permutations of the elements $\{1,2,\ldots,8\}$, and V_π is the unitary permuting the first eight factor spaces according to π.

We will now determine the asymptotic scaling of the following expression with D:

$$\mathbb{E}_C\Big\{\Pr\left(k|x,C,F(t)\right)^4\Big\} \\ = \sum_{\pi,\sigma}\langle L_8|\Phi_\pi\rangle(M^{-1})_{\pi,\sigma}\langle\Phi_\sigma|R_8(t)\rangle, \tag{B15}$$

by finding an approximation for M^{-1}. Recall that the matrix M has components $M_{\pi,\sigma} = \langle V_\pi|V_\sigma\rangle = \mathrm{Tr}[V_{\pi^{-1}}V_\sigma] = D^{l(\pi^{-1}\sigma)}$, where $l(\sigma)$ is number of cycles in the cycle decomposition of the permutation $\pi^{-1}\sigma$. It is clear that the diagonal components of M are all equal to D^8, and all other components of M are strictly $< D^8$, as the identity permutation is the only one with 8 cycles. Letting $A := \mathbb{1} - M/D^8$, we have

$$(\mathbb{1}-A)\left(\sum_{k=0}^{N}A^k\right) = \left(\sum_{k=0}^{N}A^k\right)(\mathbb{1}-A) = \mathbb{1} - A^{N+1}, \tag{B16}$$

which converges to $\mathbb{1}$ in the limit $N\to\infty$ if and only if $||A||_{\mathrm{op}} < 1$. Because A is of fixed size $8!\times 8!$ and all its elements are $< 1/D$, we have $||A||_{\mathrm{op}} < 1$ for D sufficiently large. Therefore, $(\mathbb{1}-A)^{-1} = \mathbb{1} + A + O(D^{-2})$, which implies that

$$M^{-1} = \frac{\mathbb{1}}{D^8} + O(D^{-9}). \tag{B17}$$

Next, it is not difficult to see that all components of $\langle L_8|\Phi_\pi\rangle$ are equal to 1 or δ_{xk}. Therefore, we can split the sum in (B15) as

$$\mathbb{E}_C\Big\{\Pr\left(k|x,C,F(t)\right)^4\Big\} \\ \leq \frac{1}{D^8}\sum_{\pi}\langle\Phi_\pi|R_8(t)\rangle + \sum_{\pi,\sigma}B_{\pi,\sigma}\langle\Phi_\sigma|R_8(t)\rangle, \tag{B18}$$

where we have used that $\langle L_8|\Phi_\pi\rangle \leq 1$ for all π, and $B = M^{-1} - \frac{\mathbb{1}}{D^8}$ has all components $\leq O(D^{-9})$. □

2. Expectations over GUE eigenvalues

Our next step towards proving Theorem 2 is to find the GUE average of the the spectral form factor $\mu(t)$, which appears in Lemmas 1 and 2. The joint distribution over the un-ordered energy eigenvalues $\{E_a\}_{a=1}^{D}$ of a GUE matrix is given by (see [18] Theorem 3.3.1, or [17] Chapter 4):

$$P(\{E_a\}^D) = \frac{1}{C_D}\prod_{1=a<b}^{D}(E_a-E_b)^2\exp\Big[-\frac{1}{2\sigma^2}\sum_{a=1}^{D}E_a^2\Big], \tag{B19}$$

where $C_D = (2\pi)^{D/2}\sigma^{D^2}\prod_{j=1}^{D}j!$. Integration of $P(\{E_a\}^D)$ over $D-m$ variables gives the m-point correlation function ([18] 6.1.1)

$$R_m(E_1,\ldots,E_m) = \chi\int_{-\infty}^{\infty}P(\{E_a\}^D)dx_{m+1}\cdots dx_D, \tag{B20}$$

where $\chi = D!/(D-m)!$. By [18] Theorem 5.1.4, this can be written as:

$$R_m(E_1,\ldots,E_m) = \det[K_D(E_i,E_j)]_{i,j=1,\ldots,m}, \tag{B21}$$

where $K_D(E_i,E_j) = \sum_{k=0}^{D-1}\phi_k(E_i)\phi_k(E_j)$, and the $\phi_k(x)$ are the harmonic oscillator wave-functions $\phi_j(x) = \left(2^j j!\sqrt{\pi}\right)^{-1/2}e^{-x^2/2}H_j(x)$, with $H_j(x)$ the Hermite polynomials (see [18] section 6.2).

a. *Calculation of* $\mathbb{E}_{\mathrm{spec}}\{|\mu(t)|^2\}$

Our next goal is to evaluate the expression $\Gamma(t,D) := \mathbb{E}_{\mathrm{spec}}\{|\mu(t)|^2\} = \mathbb{E}_{\mathrm{spec}}\Big\{\left|\mathrm{Tr}\left[e^{-itH}\right]\right|^2\Big\}$. Expanding out the trace and grouping terms, we have

$$\begin{aligned}\Gamma(t,D) &= \mathbb{E}_{\mathrm{spec}}\Big\{\left|\mathrm{Tr}\left[e^{-itH}\right]\right|^2\Big\} \\ &= \int_{-\infty}^{\infty}\left|\sum_{l=1}^{D}e^{itE_l}\right|^2 P(\{E_a\}^D)\prod_{j=1}^{D}dE_j \\ &= D\int_{-\infty}^{\infty}P(\{E_a\}^D)\prod_{j=1}^{D}dE_j \\ &\quad + \int_{-\infty}^{\infty}\sum_{l\neq m}^{D}e^{it(E_l-E_m)}P(\{E_a\}^D)\prod_{j=1}^{D}dE_j.\end{aligned} \tag{B22}$$

Using the invariance of the joint distribution $P(\{E_a\}^D)$ under permutations of the energies, and the definition of the 2-point correlation function, this becomes

$$\Gamma(t,D) = D + \int_{-\infty}^{\infty}e^{it(E_2-E_1)}R_2(E_1,E_2)dE_1dE_2. \tag{B23}$$

It is well known that for large D the function $K_D(E,E)$ follows the so called Wigner semi-circle law [18] (with $\sigma^2 = 1/2$):

$$K_D(E,E) \simeq \frac{1}{\pi}\sqrt{2D-E^2}\,\Theta(\sqrt{2D}-|E|), \tag{B24}$$

where $\Theta(x)$ is the Heavyside step function. For large D we then have:

$$\begin{aligned}\Gamma(t,D) &= D + 2D\frac{J_1\left(\sqrt{2D}t\right)^2}{t^2} \\ &\quad - (\sqrt{2D}-t/2)\Theta(2\sqrt{2D}-t),\end{aligned} \tag{B25}$$

where $J_1(x)$ is the first Bessel function of the first kind. It should also be noted that, the limit of the function $K_D(E_1,E_2)$ is generally taken by rescaling (often called 'unfolding' - see [24] section III.A.1) the energy level density as well as the energies by the local mean spacing [17, 18]. In taking this limit and the integrals for $\Gamma(t,D)$ we have followed the the procedure of [18] Appendices 10, and 11.

b. Equilibration time

Now that we have the GUE expectation of the spectral form factor, we can use Lemma 2, to find the full expectation over eigenvectors and eigenvalues of the outcome probabilities:

$$\begin{aligned}\mathbb{E}_{\mathrm{spec,C}}&\{\Pr(k|x,C,F(t))\}\\&=\frac{D-\frac{1}{D}\Gamma(t,D)+\delta_{xk}\left(\Gamma(t,D)-1\right)}{D^2-1},\end{aligned}\tag{B26}$$

Notice that if $\Gamma(t,D)=O(D)$, then

$$\mathbb{E}_{\mathrm{spec,C}}\{\Pr(k|x,C,F(t))\}=\frac{D+\delta_{xk}O(D)}{D^2-1}+O(D^{-2}).\tag{B27}$$

In particular, if D is large, then the GUE expectation of the probability distribution from (B27) is essentially the uniform distribution. We therefore take the *equilibration time* $t_{\mathrm{eq}}(D)$ to be defined by the condition

$$t_{\mathrm{eq}}(D):=\{T\mid\Gamma(t,D)=O(D),\forall\ t>T\}.\tag{B28}$$

It is clear that the second term in (B25) satisfies:

$$\lim_{t\to\infty}2D\frac{J_1\left(\sqrt{2D}t\right)^2}{t^2}=0.\tag{B29}$$

This shows that there exists a finite time $t_{\mathrm{eq}}(D)$ such that $\Gamma(t,D)=O(D)$ for all $t>t_{\mathrm{eq}}(D)$.

In order to get a sense of the equilibration time (keeping in mind our previous comments on GUE energy spectra), note that the condition on the equilibration time (B28) is essentially that

$$2D\frac{J_1\left(\sqrt{2D}t\right)^2}{t^2}=O(D).\tag{B30}$$

Using the fact that for $x\gg 3/4$ we can approximate $J_1(x)\simeq\sqrt{2/\pi x}\cos[x-3\pi/4]$, it the follows that the equilibration time is

$$t_{\mathrm{eq}}(D)=O(D^{-1/6}).\tag{B31}$$

c. GUE expectation of first and second power

Our next step towards a proof of Theorem 2 involves evaluating the expectation values over the spectrum that are needed to prove eqns. (3), and (4). First, from the previous section we have that there exists a finite time $t_{\mathrm{eq}}(D)$ such that $\Gamma(t,D)=O(D)$ for all $t>t_{\mathrm{eq}}(D)$. Using Lemma 2, and the above spectral expectation, we have

$$\mathbb{E}_{spec,C}\{\Pr(k|x,C,F(t))\}=\frac{D+\delta_{xk}O(D)}{D^2-1}+O(D^{-2}),\tag{B32}$$

which proves that for all $t>t_{\mathrm{eq}}(D)$, eqn. 3 holds.

Next, Lemma 1 implies that

$$\begin{aligned}&\mathbb{E}_{\mathrm{spec}}\{\mathbb{V}_C\{\Pr(k|x,C,F(t))\}\}\\&\quad=\frac{1}{D^4}\Big\{D^2-2D+4+(7-2D)\delta_{xk}\\&\quad+\Gamma(t,D)(2D\delta_{xk}-10\delta_{xk}-2)+\delta_{xk}\Gamma(2t,D)\\&\quad+2\delta_{xk}\mathbb{E}_{\mathrm{spec}}\big\{Re[\mu(t)^2\mu(-2t)]\big\}\Big\}+O(D^{-5}).\end{aligned}\tag{B33}$$

In the next section we will show that for $t>t_{\mathrm{eq}}(D)$, $\mathbb{E}_{\mathrm{spec}}\big\{Re[\mu(t)^2\mu(-2t)]\big\}=O(D)$. Using this along with $\Gamma(t,D)=O(D)$ in the above, we find

$$\mathbb{E}_{\mathrm{spec}}\{\mathbb{V}_C\{\Pr(k|x,C,F(t))\}\}=O(D^{-2}),\tag{B34}$$

which proves that for all $t>t_{\mathrm{eq}}(D)$, eqn. 4 holds.

d. Bounding $\mathbb{E}_{\mathrm{spec}}\big\{\mu(t)^2\mu(-2t)\big\}$

In order to simplify the derivation of our upper bounds, we define

$$\begin{aligned}\Delta(t,D)&:=\mathbb{E}_{\mathrm{spec}}\big\{\mu(t)^2\mu(-2t)\big\}\\&=\int_{-\infty}^{\infty}\sum_{i,j,k}e^{it(E_i+E_j-2E_k)}P(\{E_a\}^D)\prod_{l=1}^{D}dE_l,\end{aligned}\tag{B35}$$

We can expand the triple sum into three parts depending on whether i,j,k are: (i) all distinct, (ii) only two equal, or (iii) all equal, which upon integration over the remaining energies, give terms of the form:

(i) $R_3(E_1,E_2,E_3)e^{it(E_1+E_2-2E_3)}$,
(ii) $R_2(E_1,E_2)e^{2it(E_1-E_2)}+2R_2(E_1,E_2)e^{it(E_1-E_2)}$,
(iii) D.

By expanding R_3 and R_2, we find products of integrals of terms of the form $K_D(E_i,E_i)e^{it_iE_i}$, as well as $K_D(E_i,E_j)^2e^{i(t_iE_i+t_jE_j)}$, and

$$K_D(E_1,E_2)K_D(E_2,E_3)K_D(E_3,E_1)e^{it(E_1+E_2-2E_3)}.\tag{B36}$$

We will bound the magnitude of these terms in a similar fashion as was done in [11]. Note that the integral of (B36) is just

$$\mathrm{Tr}\big[Pe^{iE_1}Pe^{iE_2}Pe^{-2iE_3}\big],\tag{B37}$$

where $P=\sum_{k=0}^{D-1}|\phi_k\rangle\langle\phi_k|$ is the projector onto the D-dimensional lower-energy subspace spanned by the harmonic oscillator wave-functions, and X is the position operator. Using the Cauchy-Schwartz inequality twice on (B37), we find:

$$\begin{aligned}\big|\mathrm{Tr}\big[(Pe^{iE_1}Pe^{iE_2})Pe^{-2iE_3}\big]\big|&\leq\sqrt{\sqrt{\mathrm{Tr}[P]\,\mathrm{Tr}[P]}\mathrm{Tr}[P]}\\&\leq D.\end{aligned}\tag{B38}$$

A similar argument also shows that terms with integrands of the form $K_D(E_i, E_j)^2 e^{i(t_i E_i + t_j E_j)}$ are also bounded by D. Using these bounds, we then have

$$
\begin{aligned}
&|\Delta(t, D)| \leq 6D \\
&+ \left(\int_{-\infty}^{\infty} K_D(E,E) e^{itE}\right)^2 \left(\int_{-\infty}^{\infty} K_D(E,E) e^{-2itE}\right) \\
&+ D\left(\int_{-\infty}^{\infty} K_D(E,E) e^{-2itE}\right) + 2D\left(\int_{-\infty}^{\infty} K_D(E,E) e^{itE}\right) \\
&+ \left|\int_{-\infty}^{\infty} K_D(E,E) e^{2itE}\right|^2 + 2\left|\int_{-\infty}^{\infty} K_D(E,E) e^{itE}\right|^2 .
\end{aligned} \tag{B39}
$$

Next recall that

$$
\int_{-\infty}^{\infty} K_D(E,E) e^{itE} = \sqrt{2D}\,\frac{J_1(\sqrt{2D}\ t)}{t}, \tag{B40}
$$

which approaches 0 as $t \to \infty$. The equilibration condition (B28) requires that $\sqrt{2D} J_1(\sqrt{2D}\ t)/t = O(\sqrt{D})$ for $t > t_{\mathrm{eq}}(D)$, and if this is satisfied, then for all $t > t_{\mathrm{eq}}$, we have $|\Delta(t, D)| \leq O(D)$.

e. GUE expectation of fourth power

The final quantity that we need to prove Theorem 2 is the asymptotic scaling of the GUE average of the fourth moment of $\Pr(k|x)$. This quantity shows a concentration of measure for the outcome variance of $\Pr(k|x)$, which will allow us to conclude that individual Hamiltonians drawn from the GUE will information theoretically equilibrate with high probability. Specifically, we show that for $t > t_{\mathrm{eq}}(D)$ we have that $\mathbb{E}_{spec,C}\left\{\Pr(k|x, C, F(t))^4\right\} = O(D^{-4})$. Recalling the form of $\mathbb{E}_C\left\{\Pr(k|x, C, F(t))^4\right\}$ from Lemma 3, we see that it is sufficient to show that for $t > t_{\mathrm{eq}}(D)$, we have $\mathbb{E}_{\mathrm{spec}}\{\langle \Phi_\sigma | R_8(t)\rangle\} \leq O(D^4)$, for all permutations σ.

Calculating some explicit examples of

$$
\begin{aligned}
&\langle \Phi_\sigma | R_8(t)\rangle \\
&\quad = \sum_{\substack{a,a',b,b',\\ c,c',d,d'}} e^{it(E_a - E_{a'} + E_b - E_{b'} + E_c - E_{c'} + E_d - E_{d'})} \\
&\quad \times \langle a, a', b, b', c, c', d, d' | V_\sigma | a, a', b, b', c, c', d, d'\rangle,
\end{aligned} \tag{B41}
$$

we see that these are of the general form

$$
D^a \mu(f_1 t)\mu(g_1 t)^*, \ldots, \mu(f_4 t)\mu(g_4 t)^*, \tag{B42}
$$

where $f_j, g_j \in \{0, 1, 2, 3, 4\}$, and

$$
a + \sum_{j=1}^{4} f_j + \sum_{j=1}^{4} g_j = 8, \tag{B43}
$$

and if $f_j = 0$ then the corresponding $\mu(f_j t)$ does not appear in the product (and similarly for g_j). For example, there is a $|\mu(t)|^4$ term arising from $\sigma = \mathbb{1}$, and a D^4 term arising from $\sigma = (12)(34)(56)(78)$. The expectations $\mathbb{E}_{\mathrm{spec}}\{D^a \mu(f_1 t)\mu(g_1 t)^*, \ldots \mu(f_4 t)\mu(g_4 t)^*\}$ can then be bounded in a similar fashion to Appendix B 2 d. In particular, we can expand the sums in the product of the terms $\mu(f_j t) = \mathrm{Tr}\left[e^{i f_j t H}\right]$ into various parts depending on which indices are equal or not equal, just as was done for expression (B35). These will then give a constant D^b factor, and various combinations of integrals of m-point correlation functions. Each of the integrals with 2-point or higher order correlation functions can be bounded by D, just as was done for eqn. (B36). We will then be left with various powers of integrals of the form $\int_{-\infty}^{\infty} K_D(E,E) e^{itE}$ which as we have seen approach 0 as $t \to \infty$, and so are irrelevant for the $t > t_{eq}$ regime. The only remaining question then is power of the constant D^b factor for each term. It is not difficult to see that a power of D arises from each pairing $\mu(f_j t)\mu(g_k t)^*$ with $f_j = g_k$. For example, the expectation of the term $|\mu(t)|^4$ gives a contribution of D^2 (this is in fact the infinite time limit). From the constraint (B43), it is not difficult to see that D^4 is the highest power of D which can arise. This shows that for $t > t_{eq}$

$$
\mathbb{E}_{\mathrm{spec},C}\left\{Pr(k|i, C, F(t))^4\right\} = O(D^{-4}). \tag{B44}
$$

Putting all of the above together, we have:

Proof of Theorem 2. Theorem 2 follows directly from (B32), (B34) and (B44) for $t > t_{\mathrm{eq}}(D) = O(D^{-1/6})$, which was shown in (B31). □